

BB

USE OF O-BETA-D-GALACTOPYRANOSYL-D-XYLOSES FOR THE PREPARATION OF COMPOSITIONS AND SOLUTIONS INTENDED TO THE EVALUATION OF INTESTINAL LACTASE, AND PRODUCTION PROCESS

Número de patente: EP0819764, B1

Fecha de publicación: 1998-01-21

Inventor(es): VILLANUEVA TORREGROZA DANIEL (CO); MARTIN LOMAS MANUEL (ES); ARAGON REYES JUAN JOSE (ES); CANADA VICINAY FRANCISCO JAVIE (ES); FERNANDEZ-MAYORALAS ALVAREZ AL (ES); LOPEZ ALVAREZ ROSA (ES)

Solicitante(s): UNIV AUTONOMA (ES); CONSEJO SUPERIOR INVESTIGACION (ES)

Número de publicación: ES2100131

Número de solicitud: EP19960937342 19961108

Número(s) de prioridad: WO1996ES00208 19961108; ES19950002185 19951108

Clasificación CIP: C12Q1/34; C12P19/12; C07H3/04

Clasificación CE: C12Q1/34

Equivalentes: AU702329, AU7497796, BR9607092, CA2209927, CZ9702527, DE69619973D, DE69619973T, EA684, ES2176505T, HU9801439, JP10512457T, JP2000106898, NO973159, NZ321607, PL321257, SK100197, US5994092, WO9717464

Cited patent(s):

Resumen

There is disclosed the use of O- beta -D-galactopyranosyl-D-xyloses, particularly 2-O- beta -D-galactopyranosyl-D-xylose, and 3-O- beta -D-galactopyranosyl-D-xylose, for the preparation of compositions and solutions intended to the evaluation of the intestinal lactase, which present an affinity to the substantially enhanced lactase. There is also disclosed a process for producing O- beta -D-galactopyranosyl-D-xyloses, comprising: reacting a D-xylose and a substrate of beta -D-galactopyranoside in the presence of an enzyme beta -galactosidase, in an aqueous medium buffered to a pH comprised between 5.0 and 9.0 and at temperature between 4 and 37 DEG C; deactivating the beta -galactosidase by heating at 100 DEG C when reaching the maximum yield of formation of disaccharides detected through thin layer chromatography, and isolating the formed disaccharides through filtration in a packed column with a thinner selected between water and water/ alcohol.

Datos proporcionados por la base de datos de esp@cenet test - I2